Author Index

ABDUL-GAFFER R. *see* MACINTYRE E.A.
ADLER F.L. *see* YOSHIO T.
ADLER L.T. *see* YOSHIO T.
ADLER W.H. *see* SAXENA R.K.
AEBERSOLD R. *see* BRUDERER U.
AHMED A.E.E. & WHALEY K. Purification of a plasma protein which inhibits complement-mediated prevention of immune precipitation, 45
ALBRECHTSEN M., YEAMAN G.R. & KERR M.A. Characterization of the IgA receptor from human polymorphonuclear leucocytes, 201
ALEXANDER G.J.M. *see* NOURI-ARIA K.T.
ALKAN S.S. *see* PARAB P.B.
ANDERSON M.G. *see* NOURI-ARIA K.T.
ANDO K.-I., ISOBE K.-I., HASEGAWA T., IWAMOTO T., DING R.-N., RAHMAN J., MURO Y., YOSHIDA T., NAGASE F., KAWASHIMA K., OHASHI M. & NAKASHIMA I. Genetic and stimulator cell requirements for generation and activation of minor histocompatibility antigen-specific memory cytotoxic T-lymphocyte precursors, 661
ANGUNAWELA I.I. The effects of immunosuppression on the development of (experimental) cataract, 69
ANSTEE D.J. *see* SPRING F.A.
AOKI I. *see* YANOMA S.
ATKINSON Y.H., LOPEZ A.F., MARASCO W.A., LUCAS C.M., WONG G.G., BURNS G.F. & VADAS M.A. Recombinant human granulocyte–macrophage colony-stimulating factor (rH GM-CSF) regulates f Met-Leu-Phe receptors on human neutrophils, 519
AXELROD T.S. *see* CZITROM A.A.

BAGUERIZO A. *see* KAMADA N.
BALLINARI D. *see* DELIA D.
BANGA J.P. *see* DOBLE N.D.
BARNES P.J. *see* KROEGEL C.
BARRIOS G.A. *see* CAMPETELLA O.E.
BARTHOLOMAEUS W.N., O'DONOGHUE H., FOTI D., LAWSON C.M., SHELLAM G.R. & REED W.D. Multiple autoantibodies following cytomegalovirus infection: virus distribution and specificity of autoantibodies, 397
BATE C.A.W., TAVERNE J. & PLAYFAIR J.H.L. Malarial parasites induce TNF production by macrophages, 227
BEVERLEY P. *see* HAYWARD A.
BEVERLEY P.C.L. *see* TERRY L.A.
BHAGYAM R.C., JARRETT-ZACZEK D. & FERGUSON F.G. Activation of swine peripheral blood lymphocytes with human recombinant interleukin-2, 607
BIJSTERBOSCH M.K., MCLAUGHLIN J.B., HOLMAN M. & KLAUS G.G.B. Activation and proliferation signals in mouse B cells. IX. Protein kinase *C* activators synergize with non-mitogenic anti-immunoglobulin antibodies to drive B cells into G1, 163
BILYK N. *see* HOLT P.G.
BLASER K. *see* BRUDERER U.
BOCCI V. Roles of interferon produced in physiological conditions. A speculate review, 1
BÖCK G. *see* SCHULZ T.F.
BOND A., COOKE A. & HAY F.C. Heterogeneity in rheumatoid factor isotypes and specificities on MRL mice, 17
BOND A., HAY F.C. & COOKE A. The relationship between induced and spontaneous autoantibodies in MRL mice: the role of Ly-1 B cells, 325
BOWMAN R. *see* HOLT P.G.
BOYLSTON A. *see* HAYWARD A.
BRANDON M.R. *see* MEEUSEN E.
BRAUN D.G. *see* PARAB P.B.
BROCK J.H. *see* HUGHES A.
BROM J., SCHÖNFELD W. & KÖNIG W. Metabolism of Leukotriene B_4 by activated human polymorphonuclear granulocytes, 569
BROWN F. *see* PARRY N.R.
BROWN M.H. *see* TERRY L.A.
BROWN S. *see* PHILPOTT K.L.
BRUCE M.G. *see* LAMONT A.G.
BRUDERER U., AEBERSOLD R., BLASER K. & HEUSSER C.H. Characterization of the Group I and Group II antibodies against PC-KLH in normal and T15 idiotype-suppressed BALB/c mice, 385
BURKART V. *see* MARTIN S.
BURNS G.F. *see* ATKINSON Y.H.
BUTSCHER W.G. & FAUBERT G.M. The therapeutic action of monoclonal antibodies against a surface glycoprotein of *Giardia muris*, 175

CAMAGNA A. *see* PELOSI-TESTA E.
CAMPETELLA O.E., BARRIOS H.A. & GALASSI N.V. Suppressor T-cell population induced by Junin virus in adult mice, 407
CARAVITA T. *see* PELOSI-TESTA E.
CARREL S. *see* DE BERARDINIS P.
CARSWELL F. *see* THOMPSON S.J.
CATTORETTI G. *see* DELIA D.
CELADA A. Immune-complex inhibition of macrophage activation is not due to an interaction with the binding or processing of IFN-γ, 187
CHANDRASHEKAR R. *see* PARAB P.B.
CHANEZ P. *see* KROEGEL C.
CHIOU S.-S., YAMAUCHI K., NAKANISHI T. & OBATA H. Nature of immunological non-responsiveness to hepatitis B vaccine in healthy individuals, 545
CHOLLET-MARTIN S. *see* GOUGEROT-POCIDALO M.-A.
CHUNG K.F. *see* KROEGEL C.
COCKBURN F. *see* HUGHES A.
COLEMAN J.W. A kinetic analysis of the *in vitro* sensitization of murine peritoneal mast cells with monoclonal IgE anti-DNP antibody, 527
COOKE A. *see* BOND A.
CROMWELL O. *see* MAESTRELLI P.
CZITROM A.A., AXELROD T.S. & FERNANDES B. Granulocyte precursors are the principal cells in bone marrow that stimulate allospecific cytolytic T-lymphocyte responses, 655

DAHA M.R., GORTER A., LEIJH P.J.C., KLAR N. & VAN ES L.A. Binding and catabolism of aggregated immunoglobulins bearing C3b or iC3b by U937 cells, 375
DAHA M.R. *see* GORTER A.
DALCHAU R. *see* SPRING F.A.
DANIELS C.K. *see* SCHMUCKER D.L.
DANIELS G.L. *see* SPRING F.A.
DAVID B. *see* MECHERI S.
DAVID C.S. *see* YANOMA S.
DAVIES T.F. *see* HIROSE W.
DE BERARDINIS P., LONDEI M., CARREL S. & FELDMANN M. Regulation of clonal growth by anti-T-cell receptor antibody-directed lysis, 439
DEBS R.J. *see* VALONE F.H.
DELIA D., TRAVERSARI C., BALLINARI D., CATTORETTI G., FONTANELLA E., POLLI N. & DELLA PORTA G. Early lymphocyte activation molecule defined by the monoclonal antibody MLR-3: biochemical and functional studies, 593
DELLA PORTA G. *see* DELIA D.
DENHAM S. & HALL J.G. Studies on the adjuvant action of beryllium. III. The activity in the plasma of lymph efferent from nodes stimulated with beryllium, 341
DENT G. *see* KROEGEL C.

DEUTSCH H.F. *see* KIM H.S.
DIAMANTSTEIN T. *see* STÜNKEL K.G.
DIERICH M.P. *see* SCHULZ T.F.
DING R.-N. *see* ANDO K.-I.
DOBLE N.D., BANGA J.P., POPE R., LALOR E., KILDUFF P. & MCGREGOR A.M. Autoantibodies to the thyroid microsomal/thyroid peroxidase antigen are polyclonal and directed to several distinct antigenic sites, 23
DOINEL C. *see* LIROCHON J.
DORF M.E. *see* KUCHROO V.K.
DÖRKEN B. *see* KIESEL S.
D'SOUZA S.W. *see* TAYLOR G.M.
DUMONT F.J., PALFREE R.G.E. & FISHER P.A. The T-cell activating protein (TAP) is up-regulated by endogenous IFN-γ in activated T cells, 267

EBERSOLE J.L., STEFFEN M.J. & PAPPO J. Secretory immune responses in ageing rats. II. Phenotype distribution of lymphocytes in secretory and lymphoid tissues, 289
EBERSOLE J.L. *see* PAPPO J.
ECKELS D.D. *see* O'HEHIR R.E.
EDDLESTON A.L.W.F. *see* NOURI-ARIA K.T.
EISGI Y. & MCCULLAGH P. Acquisition of immunological self-recognition by the fetal rat, 319
VAN ES L.A. *see* DAHA M.R.
VAN ES L.A. *see* GORTER A.
ESSERY G., FELDMANN M. & LAMB J.R. Interleukin-2 can prevent and reverse antigen-induced unresponsiveness in cloned human T lymphocytes, 413
ETLINGER H.M., HEIMER E.P., TRZECIAK A., FELIX A.M. & GILLESSEN D. Assessment in mice of a synthetic peptide-based vaccine against the sporozoite stage of the human malaria parasite, *P. falciparum*, 551

FABRE J.W. *see* SPRING F.A.
FAUBERT G.M. *see* BUTSCHER W.G.
FAY M. *see* GOUGEROT-POCIDALO M.-A.
FELDMANN M. *see* DE BERARDINIS P.
FELDMANN M. *see* ESSERY G.
FELIX A.M. *see* ETLINGER H.M.
FERGUSON A. *see* LAMONT A.G.
FERGUSON A. *see* TRONCONE R.
FERGUSON F.G. *see* BHAGYAM R.C.
FERNANDES B. *see* CZITROM A.A.
FERRANTE A. & ROWAN-KELLY B. The role of antibody in immunity against experimental naegleria meningoencephalitis ('amoebic meningitis'), 241
FIEDEL B.A. Influence of tuftsin-like synthetic peptides derived from C-reactive protein (CRP) on platelet behaviour, 487
FIERA R.A. *see* NEALE M.L.
FISHER P.A. *see* DUMONT F.J.
FITZ-GIBBON L. *see* ROY S.
FONTANELLA E. *see* DELIA D.
FONTENEAU P. *see* JACHEZ B.
FORD J. *see* HOLT P.G.
FOTI D. *see* BARTHOLOMAEUS W.N.
FREW A.J. *see* O'HEHIR R.E.
FRITZ F.J. *see* PABST R.
FUJITA T., SHINKAI Y., INOUE T. & TAMURA N. Purification and characterization of decay-accelerating factor (DAF) from Raji cells, 369
FUJITA T. *see* KATAKAMI Y.
FUJITA T. *see* KOIZUMI T.
FÜST G. *see* VARGA L.

GALASSI N.V. *see* CAMPETELLA O.E.
GALLAGHER A. *see* POULTON T.A.
GARCIA-SANZ J.A., VELOTTI F., MACDONALD H.R., MASSON D., TSCHOPP J. & NABHOLZ M. Appearance of granule-associated molecules during activation of cytolytic T-lymphocyte precursors by defined stimuli, 129
GARCON N., GREGORIADIS G., TAYLOR M. & SUMMERFIELD J. Mannose-mediated targeted immunoadjuvant action of liposomes, 743
GEIST M. *see* PABST R.
GIANNELLA G. *see* PELOSI-TESTA E.
GILLESSEN D. *see* ETLINGER H.M.
GILLIS S. *see* MARTIN S.
GOOSSENS D. *see* LIROCHON J.
GORDON D.L. & RICE J.L. Opsonin-dependent and independent surface phagocytosis of *S. aureus* proceeds independently of complement and complement receptors, 709
GORRELL M.D. *see* MEEUSEN E.
GORTER A., HIEMSTRA P.S., KLAR-MOHAMAD N., VAN ES L.A. & DAHA M.R. Binding, internalization and degradation of soluble aggregates of human secretory IgA by resident rat peritoneal macrophages, 703
GORTER A., HIEMSTRA P.S., VAN DER VOORT E.A.M., VAN ES L.A. & DAHA M.R. Binding of human IgA1 to rat peritoneal macrophages, 207
GORTER A. *see* DAHA M.R.
GOUGEROT-POCIDALO M.-A., FAY M., ROCHE Y. & CHOLLET-MARTIN S. Mechanisms by which oxidative injury inhibits the proliferative response of human lymphocytes to PHA. Effect of the thiol compound 2-mercaptoethanol, 281
GREGORIADIS G. *see* GARCON N.

HAAS R. *see* KIESEL S.
HAIGH H. *see* TAYLOR G.M.
HALL J.G. Studies on the adjuvant action of beryllium. IV. The preparation of beryllium containing macromolecules that induce immunoblast responses *in vivo*, 345
HALL J.G. *see* DENHAM S.
HALLET M.-M. *see* HOUSSAINT E.
HARRIS R. *see* TAYLOR G.M.
HASEGAWA T. *see* ANDO K.-I.
HAY F.C. *see* BOND A.
HAYWARD A., BOYLSTON A. & BEVERLEY P. Lysis of CD3 hybridoma targets by cloned human CD4 lymphocytes, 87
HÉBERT J. *see* MECHERI S.
HEEG K. *see* WAGNER H.
HEIMER E.P. *see* ETLINGER H.M.
HEIN W.R., SHELTON J.N., SIMPSON-MORGAN M.W. & MORRIS B. Traffic and proliferative responses of recirculating lymphocytes in fetal calves, 621
HEIN W.R. & SUPERSAXO A. Effect of interferon-alpha-2a on the output of recirculating lymphocytes from single lymph nodes, 469
HERSCOWITZ H.B. *see* MBAWUIKE I.N.
HEUSSER C.H. *see* BRUDERER U.
HIEMSTRA P.S. *see* GORTER A.
HIROSE W. & DAVIES T.F. Generation of thyroglobulin-specific T-cell clones derived from F_1 Fisher rats, 107
HOLLY M., LIN Y.-S. & ROGERS T.J. Induction of suppressor cells by staphylococcal enterotoxin B: identification of a suppressor cell circuit in the generation of suppressor-effector cells, 643
HOLMAN M. *see* BIJSTERBOSCH M.K.
HOLT P.G., KEES U.R., SCHON-HEGRAD M.A., ROSE A., FORD J., BILYK N., BOWMAN R. & ROBINSON B.W.S. Limiting-dilution analysis of T cells extracted from solid human lung tissue: comparison of precursor frequencies for proliferative responses and lymphokine production between lung and blood T cells from individual donors, 649
HOUSSAINT E. & HALLET M.-M. Inability of adult circulating haemopoietic stem cells to sustain haemopoiesis in mouse fetal liver microenvironment,
HOWARD J.C. *see* YOUSAF N.
HUGHES A., BROCK J.H., PARROTT D.M.V. & COCKBURN F. The interaction of infant formula with macrophages: effect on phagocytic activity, relationship to expression of class II MHC antigen and survival of orally administered macrophages in the neonatal gut, 213
HUGHES C.C.W., MALE D.K. & LANTOS P.L. Adhesion of lymphocytes to cerebral microvascular cells: effects of interferon-γ, tumour necrosis factor and IL-1, 677

INOUE T. *see* FUJITA T.
ISACCHI G. *see* PELOSI-TESTA E.
ISENBERG D.A. *see* LOCNISKAR M.
ISHII N. *see* YANOMA S.
ISLAM L.N. & WILKINSON P.C. Chemotactic factor-induced polariza-

tion, receptor redistribution, and locomotion of human blood monocytes, 501
ISOBE K.-I. *see* ANDO K.-I.
IWAMOTO T. *see* ANDO K.-I.

JACHEZ B., MONTECINO-RODRIGUEZ E., FONTENEAU P. & LOOR F. Partial expression of the 1pr locus in the heterozygous state: presence of autoantibodies, 31
JARRETT-ZACZEK D. *see* BHAGYAM R.C.
JENKINSON E.J. *see* PARKIN I.G.
JOBIN M. *see* MECHERI S.
JODOUIN C.-A. *see* TALOR E.
JOHNSON J.P. *see* SCHULZ T.F.
JUDSON P.A. *see* SPRING F.A.

KAMADA N., SUMIMOTO R., BAGUERIZO A., YOSHIMATSU A., TERAMOTO K. & YAMAGUCHI A. Mechanisms of transplantation tolerance induced by liver grafting in rats: involvement of serum factors in clonal deletion, 315
KATAKAMI N. *see* KATAKAMI Y.
KATAKAMI Y., NAKAO Y., KOIZUMI T., KATAKAMI N., OGAWA R. & FUJITA T. Regulation of tumour necrosis factor production by mouse peritoneal macrophages: the role of cellular cyclic AMP, 719
KATAKAMI Y. *see* KOIZUMI T.
KAWADE Y. *see* WATANABE Y.
KAWANO Y.-I. *see* TANIGUCHI K.
KAWASHIMA K. *see* ANDO K.-I.
KAY A.B. *see* MAESTRELLI P.
KAY A.B. *see* O'HEHIR R.E.
KEES U.R. *see* HOLT P.G.
KERR M.A. *see* ALBRECHTSEN M.
KIERSZENBAUM F. *see* MOLINA H.A.
KIESEL S., PEZZUTTO A., HAAS R., MOLDENHAUER G. & DÖRKEN B. Functional evaluation of CD19- and CD22-negative variants of B-lymphoid cell lines, 445
KILDUFF P. *see* DOBLE N.D.
KIM H.S. & DEUTSCH H.F. Variable-region sequence homologies of human κ-chains with idiotypic serologic similarities, 573
KLAR N. *see* DAHA M.R.
KLAR-MOHAMAD N. *see* GORTER A.
KLAUS G.G.B. *see* BIJSTERBOSCH M.K.
KOIZUMI T., NAKAO Y., MATSUI T., KATAKAMI Y., MIHARA K., TAKAHASHI R., MAEDA S., SUGIYAMA T. & FUJITA T. *C-myc* expression in lymphocytes of MRL/MP-1pr mice activated by A23187 and TPA, 93
KOIZUMI T. *see* KATAKAMI Y.
KOLB H. *see* MARTIN S.
KOMATSU M., YAMAMOTO K., NAKANO Y., NAKAZAWA M., OZAWA A., MIKAMI H., TOMITA M. & MIGITA S. Hereditary C3 hypocomplementemia in the rabbit, 363
König W. *see* BROM J.
KÖNIG W. *see* RAULF M.
KROEGEL C., YUKAWA T., DENT G., CHANEZ P., CHUNG K.F. & BARNES P.J. Platelet-activating factor induces eosinophil peroxidase release from purified human eosinophils, 559
KUCHROO V.K., NOMA T., MINAMI M. & DORF M.E. Down-regulation of suppressor cell induction, 633

LACKIE A.M. & VASTA G.R. The role of galactosyl-binding lectin in the cellular immune response of the cockroach *Periplaneta americana* (Dictyoptera), 353
LALOR E. *see* DOBLE N.D.
LAMB J.R. *see* ESSERY G.
LAMB J.R. *see* LARCHÉ M.
LAMB J.R. *see* O'HEHIR R.E.
LAMONT A.G., BRUCE M.G., WATRET K.C. & FERGUSON A. Suppression of established DTH response to ovalbumin in mice by feeding antigen after immunization, 135
LAMONT A.G. *see* MOWAT A.McI.
LANTOS P.L. *see* HUGHES C.C.W.
LAPEYRE J. *see* MECHERI S.
LARCHÉ M., LAMB J.R. & RITTER M.A. A novel T-lymphocyte molecule that may function in the induction of self-tolerance and MHC-restriction within the human thymic microenvironment, 101
LAWSON C.M. *see* BARTHOLOMAEUS W.N.
LEIJH P.J.C. *see* DAHA M.R.
LESLIE G.A. *see* STEELE M.G.
LEVITT D. Naturally occurring mouse antibodies against T-cell-secreted chondroitin sulphate proteoglycan, 587
LIE W.-R., ROTHSCHILD M.F. & WARNER C.M. Preparation and characterization of murine monoclonal antibodies to swine lymphocyte antigens, 599
LIN Y.-S. *see* HOLLY M.
LINCH D.C. *see* MACINTYRE E.A.
LIROCHON J., DOINEL C., ROUGER P., GOOSSENS D. & SALMON C. Biochemical identification and characterization of the G antigen in the human blood group Rh, 337
LOCNISKAR M., ZUMLA A., MUDD D.W., ISENBERG D.A., WILLIAMS W. & MCADAM K.P.W.J. Human monoclonal antibodies to phenolic glycolipid-1 derived from patients with leprosy, and production of specific anti-idiotypes, 245
LONDEI M. *see* DE BERARDINIS P.
LOOR F. *see* JACHEZ B.
LOPEZ A.F. *see* ATKINSON Y.H.
LUCAS C.M. *see* ATKINSON Y.H.

MCADAM K.P.W.J. *see* LOCNISKAR M.
MCCULLAGH P. *see* EISHI Y.
MACDONALD H.R. *see* GARCIA-SANZ J.A.
MCGEE D.W. & MCMURRAY D.N. Protein malnutrition reduces the IgA immune response to oral antigen by altering B-cell and suppressor T-cell functions, 697
MCGREGOR A.M. *see* DOBLE N.D.
MACINTYRE E.A., TATHAM P.E.R., ABDUL-GAFFAR R. & LINCH D.C. The effects of pertussis toxin on human T lymphocytes, 427
MCLAUGHLIN J.B. *see* BIJSTERBOSCH M.K.
MCMASTER W.R. *see* WILLIAMS L.D.
MCMURRAY D.N. *see* MCGEE D.W.
MAEDA S. *see* KOIZUMI T.
MAESTRELLI P., TSAI J.-J., CROMWELL O. & KAY A.B. The identification and partial characterization of a human mononuclear cell-derived neutrophil chemotatic factor apparently distinct from IL-1, IL-2, GM-CSF, TNF and IFN-γ, 219
MAGRIN S. *see* NOURI-ARIA K.T.
MALE D.K. *see* HUGHES C.C.W.
MALLINSON G. *see* SPRING F.A.
MARASCO W.A. *see* ATKINSON Y.H.
MARTIN S., MARUTA K., BURKART V., GILLIS S. & KOLB H. IL-1 and IFN-γ increase vascular permeability, 301
MARUTA K. *see* MARTIN S.
MASSON D. *see* GARCIA-SANZ J.A.
MATSUI T. *see* KOIZUMI T.
MATTHEWS N. *see* NEALE M.L.
MBAWUIKE I.N. & HERSCOWITZ H.B. Relationship between ineffective antigen presentation by murine alveolar macrophages and their immunosuppressive function, 61
MECHERI S., MOURAD W., LAPEYRE J., JOBIN M., DAVID B. & HÉBERT J. Demonstration of idiotypes expressed on basophil-bound IgE antibodies by using anti-idiotype-induced histamine release in grass pollen allergic patients, 11
MEEUSEN E., GORRELL M.D. & BRANDON M.R. Presence of a distinct $CD8^+$ and $CD5^-$ leucocyte subpopulation in the sheep liver, 615
MELLOR A.L. *see* PHILPOTT K.L.
MIGITA S. *see* KOMATSU M.
MIHARA K. *see* KOIZUMI T.
MIKAMI H. *see* KOMATSU M.
MINAMI M. *see* KUCHROO V.K.
MITTERER M. *see* SCHULZ T.F.
MOLDENHAUER G. *see* KIESEL S.
MOLINA H.A. & KIERSZENBAUM F. Immunohistochemical detection of deposits of eosinophil-derived neurotoxin and eosinphil peroxidase in the myocardium of patients with Chagas' disease, 725
MONTECINO-RODRIGUEZ E. *see* JACHEZ B.
MONTESORO E. *see* PELOSI-TESTA E.
MOORE T.C., SPRUCK C.H. & SAID S.I. Depression of lymphocyte traffic in sheep by vasoactive intestinal peptide (VIP), 475
MORRIS B. *see* HEIN W.R.
MOURAD W. *see* MECHERI S.
MOUZAKI A. *see* STÜNKEL K.G.

MOWAT A. MCI., LAMONT A.G. & PARROTT D.M.V. Suppressor T cells, antigen-presenting cells and the role of I-J restriction in oral tolerance to ovalbumin, 141
MUDD D.W. *see* LOCNISKAR M.
MURO Y. *see* ANDO K.-I.

NABHOLZ M. *see* GARCIA-SANZ J.A.
NAGASE F. *see* ANDO K.-I.
NAKANISHI T. *see* CHIOU S.-S.
NAKANO Y. Non-specific regulatory mechanism of contact sensitivity: the requirement of intermediate cells for non-specific suppressor factor (NSF) activity, 261
NAKANO Y. *see* KOMATSU M.
NAKAO Y. *see* KATAKAMI Y.
NAKAO Y. *see* KOIZUMI T.
NAKASHIMA I. *see* ANDO K.-I.
NAKAZAWA M. *see* KOMATSU M.
NAYSMITH J.D. *see* THOMPSON S.J.
NEALE M.L., FIERA R.A. & MATTHEWS N. Involvement of phospholipase A2 activation in tumour cell killing by tumour necrosis factor, 81
NOMA T, *see* KUCHROO V.K.
NOMOTO K. *see* TANIGUCHI K.
NOURI-ARIA K.T., MAGRIN S., ALEXANDER G.J.M., ANDERSON M.G., WILLIAMS R. & EDLESTON A.L.W.F. Abnormal T-cell activation in chronic hepatitis B viral infection: a consequence of monocyte dysfunction? 733

OBATA H. *see* CHIOU S.-S.
O'DONOGHUE H. *see* BARTHOLOMAEUS W.N.
OGAWA R. *see* KATAKAMI Y.
OH S.K., ROSS S., WALKER J. & ZEISEL S. Role of a SER immune suppressor in immune surveillance, 73
OHASHI M. *see* ANDO K.-I.
O'HEHIR R.E., ECKELS D.D., FREW A.J., KAY A.B. & LAMB J.R. MHC class II restriction specificity of cloned human T lymphocytes reactive with *Dermatophagoides farinae* (house dust mite), 627
OKUDA K. *see* YANOMA S.
O'NEILL H.C. & PARISH C.R. Monosaccharide inhibition of cytotoxic T-cell function: demonstration of clone-specific effects, 181
OWEN J.J.T. *see* PARKIN I.G.
OZAWA A. *see* KOMATSU M.

PABST R., GEIST M., ROTHKÖTTER H.J. & FRITZ F.J. Postnatal development and lymphocyte production of jejunal and ileal Peyer's patches in normal and gnotobiotic pigs, 539
PALFREE R.G.E. *see* DUMONT F.J.
PARAB P.B., RAJASEKARIAH G.R., CHANDRASHEKAR R., ALKAN S.S., BRAUN D.G. & SUBRAHMANYAM D. Characterization of a monoclonal antibody against infective larvae of *Brugia malyai*, 169
PARISH C.R. *see* O'NEILL H.C.
PARKIN I.G., OWEN J.J.T. & JENKINSON E.J. Proliferation of thymocytes in relation to T-cell receptor β-chain expression, 97
PARROTT D.M.V. *see* HUGHES A.
PARROTT D.M.V. *see* MOWAT A.MCI.
PARRY N.T., SYRED A., ROWLANDS D.J. & BROWN F. A high proportion of anti-peptide antibodies recognize foot-and-mouse disease virus particles, 567
PARSONS S.F. *see* SPRING F.A.
PATTERSON R. *see* PRUZANSKY J.J.
PAPPO J., EBERSOLE J.L. & TAUBMAN M.A. Phenotype of mononuclear leucocytes resident in rat major salivary and lacrimal glands, 295
PAPPO J. *see* EBERSOLE J.L.
PELOSI-TESTA E., SAMOGGIA P., GIANNELLA G., MONTESORO E., CARAVITA T., SALVO G., CAMAGNA A., ISACCHI G., TESTA U. & PESCHLE C. Mechanisms underlying T-lymphocyte activation: mitogen initiates and IL-2 amplifies the expression of transferrin receptors via intracellular iron level, 273
PESCHLE C. *see* PELOSI-TESTA E.
PEZZUTTO A. *see* KIESEL S.
PHILIP R. *see* VALONE F.H.
PHILPOTT K.L., RASTAN S., BROWN S. & MELLOR A.L. Expression of H-2 class I genes in murine extra-embryonic tissues, 479
PLAYFAIR J.H.L. *see* BATE C.A.W.
POLLI N. *see* DELIA D.
POPE R. *see* DOBLE N.D.
POTTS R.C. *see* POULTON T.A.
POULIN L. *see* ROY S.
POULTON T.A., GALLAGHER A., POTTS R.C. & SWANSON BECK J. Changes in activation markers and cell membrane receptors on human peripheral blood T lymphocytes during cell cycle progression after PHA stimulation, 419
PRUZANSKY J.J. & PATTERSON R. Limiting concentrations of human basophil-bound IgE antibody required for histamine release, 307

RAHMAN J. *see* ANDO K.-I.
RAJASEKARIAH G.R. *see* PARAB P.B.
RASTAN S. *see* PHILPOTT K.L.
RAULF M. & KÖNIG W. Modulation of leukotriene release from human polymorphonuclear leucocytes by PMA and arachidonic acid, 51
REED W.D. *see* BARTHOLOMAEUS W.N.
RICE J.L. *see* GORDON D.L.
RICHTER M. *see* TALOR E.
RITTER M.A. *see* LARCHÉ M.
ROBINSON B.W.S. *see* HOLT P.G.
ROCHE Y. *see* GOUGEROT-POCIDALO M.-A.
ROGERS T.J. *see* HOLLY M.
ROSE A. *see* HOLT P.G.
ROSS S. *see* OH S.K.
ROTHKÖTTER H.J. *see* PABST R.
ROTHSCHILD M.F. *see* LIE W.-R.
ROUGER P. *see* LIROCHON J.
ROWAN-KELLY B. *see* FERRANTE A.
ROWLANDS D.J. *see* PARRY N.R.
ROY S., FITZ-GIBBON L., POULIN L. & WAINBERG M.A. Infection of human monocytes/macrophages by HIV-1: effect on secretion of IL-1 activity, 233

SAID S.I. *see* MOORE T.C.
SAITO K. *see* YOSHIO T.
SALMON C. *see* LIROCHON J.
SALVO G. *see* PELOSI-TESTA E.
SAMOGGIA P. *see* PELOSI-TESTA E.
SAXENA R.K., SAXENA Q.B. & ADLER W.H. Lectin-induced cytotoxic activity in spleen cells from young and old mice. Age-related changes in types of effector cells, lymphokine production and response, 457
SAXENA Q.B. *see* SAXENA R.K.
SCHLUMBERGER H.D. *see* STÜNKEL K.G.
SCHMUCKER D.L., DANIELS C.K., WANG R.K. & SMITH K. Mucosal immune response to cholera toxin in ageing rats. I. Antibody and antibody-containing cell response, 691
SCHÖNFELD W. *see* BROM J.
SCHON-HEGRAD M.A. *see* HOLT P.G.
SCHULZ T.F., VOGETSEDER W., MITTERER M., BÖCK G., JOHNSON J.P. & DIERICH M.P. Individual epitopes of an 85,000 MW membrane adherence molecule are variably expressed on cells of different lineage, 581
SHELTON J.N. *see* HEIN W.R.
SHELLAM G.R. *see* BARTHOLOMAEUS W.N.
SHINKAI Y. *see* FUJITA T.
SILVENNOINEN O. Purified human NK cells do not function as accessory cells in T-cell proliferative responses, 495
SIMPSON-MORGAN M.W. *see* HEIN W.R.
SMITH K. *see* SCHMUCKER D.L.
SPRING F.A., DALCHAU R., DANIELS G.L., MALLINSON G., JUDSON P.A., PARSONS S.F., FABRE J.W. & ANSTEE D.J. The In^a and In^b blood group antigens are located on a glycoprotein of 80,000 MW (the CDw44 glycoprotein) whose expression is influenced by the *In (Lu)* gene, 37
SPRUCK C.H. *see* MOORE T.C.
STEELE M.G. & LESLIE G.A. IgD antibodies to DNP are specifically induced in ascitic fluid after intraperitoneal immunization, 391
STEFFEN M.J. *see* EBERSOLE J.L.
STÜNKEL K.G., THEISEN P., MOUZAKI A., DIAMANTSTEIN T. & SCHLUMBERGER H.D. Monitoring of interleukin-2 receptor (IL-2R) expression *in vivo* and studies on an IL-2R-directed immunosuppressive therapy of active and adoptive adjuvant-induced arthritis in rats, 683
SUBRAHMANYAM D. *see* PARAB P.B.
SUGIYAMA T. *see* KOIZUMI T.

SUMIMOTO R. *see* KAMADA N.
SUMMERFIELD J. *see* GARCON N.
SUPERSAXO A. *see* HEIN W.R.
SWAIN S.L. *see* TONKONOGY S.L.
SWANSON BECK J. *see* POULTON T.A.
SYRED A. *see* PARRY N.R.

TAKAHASHI R. *see* KOIZUMI T.
TALOR E., JODOUIN C.-A. & RICHTER M. Cells involved in the immune response. XXXVI. The thymic antigen-specific suppressor cell in the immunized rabbit is a T cell with receptors for FcG and the antigen and it reacts, via a secreted suppressor factor, directly on the immune splenic AFC B cell to inhibit antibody secretion, 253
TAMURA N. *see* FUJITA T.
TANI K. *see* YANOMA S.
TANIGUCHI K., KAWANO Y.-I. & NOMOTO K. AK cells were developed from NK cells during *in vitro* culture of allogeneic or F_1 anti-parental stimulation: functional conversion in recognizing H-2 expression of target cells accompanied by phenotypical conversion, 669
TATHAM P.E.R. *see* MACINTYRE E.A.
TAUBMAN M.A. *see* PAPPO J.
TAVERNE J. *see* BATE C.A.W.
TAYLOR G.M., HAIGH H., WILLIAMS A., D'SOUZA S.W. & HARRIS R. Down's syndrome lymphoid cell lines exhibit increased adhesion due to the over-expression of lymphocyte function-associated antigen (LFA-1), 451
TAYLOR M. *see* GARCON N.
TEH H.-S. *see* WILLIAMS L.D.
TERAMOTO K. *see* KAMADA N.
TERRY L.A., BROWN M.H. & BEVERLEY P.C.L. The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45, 331
TESTA U. *see* PELOSI-TESTA E.
THEISEN P. *see* STÜNKEL K.G.
THIRY E. *see* VARGA L.
THOMPSON S.J., WHITLEY H.J., NAYSMITH J.D. & CARSWELL F. IgE antibodies to *D. pteronyssinus* in atopic patients, 311
TOMITA M. *see* KOMATSU M.
TONKONOGY S.L. & SWAIN S.L. Stimulation of gut-associated lymphoid cells by IL-4 and B-cell growth factor II, 155
TRAVERSARI C. *see* DELIA D.
TRONCONE R. & FERGUSON A. In mice, gluten in maternal diet primes systemic immune responses to gliadin in offspring, 533
TRZECIAK A. *see* ETLINGER H.M.
TSAI J.-J. *see* MAESTRELLI P.
TSCHOPP J. *see* GARCIA-SANZ J.A.

VADAS M.A. *see* ATKINSON Y.H.
VALONE F.H., PHILIP R. & DEBS R.J. Enhanced human monocyte cytotoxicity by platelet-activating factor, 715
VARGA L., THIRY E. & FÜST G. BSA–anti-BSA immune complexes formed in the presence of human complement do not bind to autologous red blood cells, 381
VASTA G.R. *see* LACKIE A.M.
VELOTTI F. *see* GARCIA-SANZ J.A.
VOGETSEDER W. *see* SCHULZ T.F.
VAN DER VOORT E.A.M. *see* GORTER A.

WAGNER H. & HEEG K. Two distinct signals regulate induction of IL-2 responsiveness in $CD8^+$ murine T cells, 433
WAINBERG M.A. *see* ROY S.
WALKER J. *see* OH S.K.
WANG R.K. *see* SCHMUCKER D.L.
WARNER C.M. *see* LIE W.-R.
WATANABE Y. & KAWADE Y. Suppressive effects of mouse interferon-β on gene expression occurring in concanavalin A-stimulated mouse spleen cells, 739
WATRET K.C. *see* LAMONT A.G.
WHALEY K. *see* AHMED A.E.E.
WHITLEY H.J. *see* THOMPSON S.J.
WILLIAMS A. *see* TAYLOR G.M.
WILLIAMS B.D. *see* YOUSAF N.
WILLIAMS L.D., MCMASTER W.R. & TEH H.-S. Antigen-independent activation of cytotoxic T cells by lymphokines, 121
WILLIAMS R. *see* NOURI-ARIA K.T.
WILLIAMS W. *see* LOCNISKAR M.
WILKINSON P.C. *see* ISLAM L.N.
WONG G.G. *see* ATKINSON Y.H.

YAMAGUCHI A. *see* KAMADA N.
YAMAMOTO K. *see* KOMATSU M.
YAMAUCHI K. *see* CHIOU S.-S.
YANOMA S., AOKI I., ISHII N., TANI K., DAVID C.S. & OKUDA K. The role of autoreactive T-cell hybridomas from autoimmune model mice, 113
YUKAWA T. *see* KROEGEL C.
YEAMAN G.R. *see* ALBRECHTSEN M.
YOSHIDA T. *see* ANDO K.-I.
YOSHIMATSU A. *see* KAMADA N.
YOSHIO T., SAITO K., ADLER F.L. & ADLER L.T. A role for mature B cells in bone marrow transplantation, 147
YOUSAF N., HOWARD J.C. & WILLIAMS B.D. Studies in the rat of antibody-sensitized and *N*-ethylmaleimide-treated erythrocyte clearance by the liver: effects of immune complex infusion and complement activation, 193

ZEISEL S. *see* OH S.K.
ZUMLA A. *see* LOCNISKAR M.

Subject Index

Adherence molecule, 85,000 MW:
  variable epitope expression, 581
Adhesion, by trisomy 21 cells:
  expression of LFA-1, 451
Adhesion of lymphocytes, to cerebral microvascular cells:
  effects of IFN-γ, TNF and IL-1, 677
Aggregated immunoglobulins:
  binding and catabolism, 375
Allospecific cytolytic T lymphocytes:
  stimulus by granulocyte precursors, 655
Alveolar macrophages:
  mechanisms of suppression, 61
Antibody:
  role in immunity to amoebic meningitis, 241
Anti-idiotypes:
  on human antibodies to leprosy phenolic glycolipid-1, 245
Autoantibodies following cytomegalovirus infection, 397
Autoantibodies, induced and spontaneous, in MRL mice:
  role of Ly-1 B cells, 325
Autoantibodies to thyroid microsomal/thyroid peroxidase antigen, 23
Autoreactive T-cell hybridomas:
  from autoimmune mice, 113

B cells:
  in bone marrow transplantation, 147
B cells, mouse:
  activation and proliferation signals, 163
Beryllium:
  adjuvant action, 341
  adjuvant action, induction of immunoblast responses *in vivo*, 345
Blood group antigens:
  In^a and In^b, on the CDw44 glycoprotein, 87
B-lymphoid cell lines:
  CD19- and CD22-negative variants, 445
Bone marrow transplantation:
  role for B cells in, 147
Brugia malayi, infective larvae:
  monoclonal antibody against, 169

C3 hypocomplementemia:
  in rabbits, 363
Cataract, experimental:
  effects of immunosuppression, 69
CD4 lymphocytes, cloned, human:
  lysis of CD3 hybridomas, 87
$CD8^+$ $CD5^-$ leucocyte subpopulation in sheep liver, 615
CD8 T-cell activation signals, 433
CD19- and CD22-negative variant B-cell lines, 445
Chemotactic factor, neutrophil, human:
  mononuclear cell-derived, 219
Cholera toxin,
  mucosal immune reponse to, 691
Class I, expression:
  in murine extra-embryonic tissue, 479
Clonal growth:
  regulation by anti-T-cell receptor antibody-directed lysis, 439
C-myc expression in MRL/MP-1pr mouse lymphocytes, 93
Complement:
  and surface phagocytosis, 709
Complement-mediated inhibition of precipitation:
  purification of plasma protein inhibitor, 45
C-reactive protein, tuftsin-like peptides from:
  influence on platelets, 487
Cytolytic T-cell precursors:
  granule-associated molecules, during activation, 129
Cytomegalovirus infection:
  and multiple autoantibody production, 397
Cytotoxic T-cell function:
  monosaccharide inhibition, 181

Decay-accelerating factor (DAF):
  Raji cells, purification and characterization, 369
DTH response:
  suppression by feeding antigen, 135

Eosinophil proteins:
  in chagastic myocardial lesions, 725
Epitope expression, variable:
  of 85,000 MW adherence molecule, 581

Foot-and-mouth disease virus:
  recognized by anti-peptide antibodies, 567

Galactosyl-binding lectin:
  role in cellular immune response of the cockroach, 353
GALT, B cells:
  lymphokine stimulation of, 155
Giardia muris:
  therapeutic action of monoclonal antibody against a surface glycoprotein, 175
Gluten, in maternal diet:
  primes responses to gliadin in offspring, 533
Granulocyte–macrophge colony-stimulating factor:
  human, recombinant, effect on neutrophils, 519
Group I and II anti-PC-KLH antibodies:
  in T15 idiotype-suppressed mice, 385

Haemopoietic stem cell development:
  adult, in fetal microenvironment, 463
Hepatitis B infection:
  chronic, abnormal T-cell activation, 733
Hepatitis B vaccine:
  non-responsiveness to, 545
HIV-1:
  infection of human macrophages/monocytes, 233
Human leucocyte-common antigen CD45:
  recognition of 180,000 MW component by monoclonal antibody, 331

Idiotypes:
  expressed on basphophil-bound IgE, 11
IFN-alpha-2a:
  effect on lymphocyte recirculation, 469
IFN-β:
  suppressive effects on spleen cells, 739
IFN, roles of:
  produced in physiological conditions, 1
IgA receptor:
  human PMN, characterization of, 201
IgA1, human:
  binding to rat peritoneal macrophages, 207
IgD anti-DNP antibodies, in ascitic fluid, 391
IgE antibodies:
  to *D. pteronyssinus* in atopic patients, 311
IgE, basophil bound:
  limiting concentration for histamine release, 307

IL-1 secretion by human monocytes/macrophages:
effect of HIV-1 infection, 233
IL-2:
effect on unresponsiveness in cloned human T cells, 413
IL-2 receptor:
in adjuvant arthritis, 683
IL-2, recombinant:
activation of swine lymphocytes, 607
Immune complex:
inhibition of macrophage activation, 187
Immune complexes, in presence of human complement,
do not bind autologous red blood cells, 381
Immune surveillance:
role of immune suppressor, 73
Immunosuppression:
effects on development of experimental cataract, 69

Lectin-induced cytotoxic activity:
in spleen cells, 457
Leukotriene B_4 metabolism:
by activated human polymorphs, 569
Leukotriene release:
modification by PMA and arachidonic acid, 51
Liposomes:
targeted adjuvant action, 743
1pr locus:
partial expression, autoantibodies, 31
Ly-1 B cells:
role in autoantibody production in MRL mice, 325
Lymphocyte activation markers, 419
Lymphocyte activation molecule:
biochemical and functional studies, 593
Lymphokine, IL-4 and BCGF-II:
stimulation of GALT, 155

Macrophage activation:
immune complex inhibition of, 187
Macrophages:
effect of infant formula on phagocytic activity, MHC class II expression and survival, 213
handling of secretory IgA aggregates, 703
Mast cells, murine peritoneal:
in vitro sensitization, 527
MHC restriction, class II:
of cloned human T cells, 627
Minor histocompatibility antigen:
specific cytotoxic T-cell precursors, 661
Monocyte polarization and locomotion:
chemotactic factor-induced, 501
Mononuclear leucocytes, phenotype of:
in rat salivary and lacrimal glands, 295
Mononuclear phagocyte system, hepatic, function, 193
MRL mice:
heterogeneity of rheumatoid factor, 17
MRL/MP-1pr mouse lymphocytes:
c-myc expression, 93
Mucosal immune response to cholera toxin, 691

Natural anti-proteoglycan antibodies, 587
NK cells develop into AK cells *in vitro*, 669
NK cells, human:
in T-cell proliferative responses, 495
Non-specific suppressor factor:
in contact sensitivity, 261

Oral tolerance:
role of suppressor T cells, antigen-presenting cells and I-J restriction, 141
Oxidative injury:
immune, in human lymphocytes, 281

Pertussis toxin:
effects on human T lymphocytes, 427
Peyer's patches, in pigs, development of, 539
Phenolic glycolipid-I:
from leprosy patients, monoclonal antibodies to, anti-idiotypes, 245
Phospholipase A2:
activation, in tumour cell killing by tumour necrosis factor, 81
Platelet-activating factor:
enhances human monocyte cytotoxicity, 715
induces release of eosinophil peroxidase, 559
Protein malnutrition:
effect on responses to oral antigen, 697

Recirculating lymphocytes:
in fetal calves, 621
Rheumatoid factor:
heterogeneity of, in MRL mice, 17
Rh (G) antigen:
biochemical identification and characterization, 337

Secretory IgA aggregates:
handling by peritoneal macrophages, 703
Secretory immune response, in ageing rats:
phenotype distribution of lymphocytes, 289
Self-recognition:
acquisition of, in fetal rat, 319
Suppressor cell circuit:
induced by staphylococcal enterotoxin B, 643
Suppressor cell induction:
down-regulation by TsF_2, 633
Suppressor T cells:
induced by Junin virus, 407
Surface phagocytosis:
and complement, 709
Swine lymphocyte antigens:
monoclonal antibodies to, 599
Synthetic peptide vaccine:
against sporozoites of *P. falciparum*, 551

T-cell activating protein (TAP):
regulation by endogenous IFN-γ, 267
T-cell activation:
abnormal, in chronic hepatitis B, 733
by lymphokines, 121
T-cell clones, rat:
specific for thyroglobulin, 107
T-cell receptor β-chain:
and proliferation of thymocytes, 97
T cells, human:
comparison of proliferation and lymphokine production by lung and blood T cells, 649
T-lymphocyte activation:
expression of transferrin receptors, 273
T-lymphocyte molecule:
involved in self tolerance and MHC restriction, 101
Thymic antigen-specific suppressor cells, rabbit,
factors, 253
Thymocyte proliferation:
and T-cell receptor β-chain expression, 97
Thyroglobulin:
specific T-cell clones, rat, 107
TNF by macrophages:
induced by malarial parasites, 227
Transplantation tolerance:
induced by liver grafting, role of serum factors, 315
Tuftsin-like peptides, from CRP:
effect on platelets, 487
Tumour necrosis factor:
phospholipase A2 activation in tumour cell killing, 81
regulation of production, 719

Vascular permeability:
increased by IL-1 and IFN-γ, 301
Vasoactive intestinal peptide:
depression of lymphocyte traffic in sheep, 475
V-region epitopes of human κ-chains, 573